Journal of MOLECULAR BIOLOGY

122

EDITOR-IN-CHIEF
J. C. KENDREW

ASSISTANT EDITOR
S. BRENNER

EDITORIAL BOARD
G. Attardi
S. Bernhard
J. E. Darnell
G. von Ehrenstein
G. Felsenfeld
J. Hurwitz
H. E. Huxley
A. D. Kaiser
A. Klug
O. Maaløe
A. Novick
G. E. Palade
D. C. Phillips
L. Stryer
C. Thomas
A. Tissières
J. Wyman

LIBRARY USE ONLY

UNIVERSITY OF HAWAII
LIBRARY

VOLUME 43
14 July 1969 to 14 August 1969

ACADEMIC PRESS
LONDON and NEW YORK

301
173

Copyright © 1969, by Academic Press Inc. (London) Ltd.

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 43

Number 1, 14 July 1969

Cooper, S., Cell Division and DNA Replication following a Shift to a Richer Medium 1

Rychlík, I., Černá, J., Chládek, S., Žemlička, J. and Haladová, Z., Substrate Specificity of Ribosomal Peptidyl Transferase: 2′(3′)-*O*-Aminoacyl Nucleosides as Acceptors of the Peptide Chain on the Amino Acid Site . 13

Wang, J. C., Variation of the Average Rotation Angle of the DNA Helix and the Superhelical Turns of Covalently Closed Cyclic λ DNA . . . 25

Wallace, S. S., Erlanger, B. F. and Beiser, S. M., Antibodies Specific for Purine and Pyrimidine Nucleosides: Effect on *in vitro* Priming Ability of DNA 41

Imamoto, F., Intragenic Initiations of Transcription of the Tryptophan Operon in *Escherichia coli* following Dinitrophenol Treatment without Tryptophan 51

Hawthorne, D. C., Identification of Nonsense Codons in Yeast . . . 71

Sakore, T. D. and Sobell, H. M., Crystal and Molecular Structure of a Hydrogen-bonded Complex containing Adenine and Hypoxanthine Derivatives: 9-Ethyl-8-bromoadenine–9-Ethyl-8-bromohypoxanthine . . . 77

Mueller, K. and Bremer, H., Heterogeneous Initiation and Termination of Enzymically Synthesized Ribonucleic Acid 89

Appendix—Termination and Length of RNA Molecules at Different Reaction Times. 105

Bremer, H. and Mueller, K., Substrate Requirements for Initiation and Chain Growth of Enzymically Synthesized RNA 109

Hunt, T., Hunter, T. and Munro, A., Control of Haemoglobin Synthesis: Rate of Translation of the Messenger RNA for the α and β chains . . . 123

Martin, T. E., Rolleston, F. S., Low, R. B. and Wool, I. G., Dissociation and Reassociation of Skeletal Muscle Ribosomes 135

Martin, T. E. and Wool, I. G., Active Hybrid 80 s Particles formed from Subunits of Rat, Rabbit and Protozoan (*Tetrahymena pyriformis*) Ribosomes 151

Soffer, R. L. and Horinishi, H., Enzymic Modification of Proteins. I. General Characteristics of the Arginine-transfer Reaction in Rabbit Liver Cytoplasm 163

Model, P., Webster, R. E. and Zinder, N. D., The UGA Codon *in vitro*: Chain Termination and Suppression 177

Hohn, T., Role of RNA in the Assembly Process of Bacteriophage fr . . 191

Reznikoff, W. S., Miller, J. H., Scaife, J. G. and Beckwith, J. R., A Mechanism for Repressor Action 201

Letters to the Editor

Reznikoff, W. S. and Beckwith, J. R., Genetic Evidence that the Operator Locus is distinct from the *z* gene in the *lac* Operon of *Escherichia coli* 215

Bhorjee, J. S., Fowler, A. V. and Zabin, I., Biochemical Evidence that the Operator Locus is distinct from the *z* Gene in the *lac* Operon of *Escherichia coli* 219

Humphrey, R. L., Avey, H. P., Becka, L. N., Poljak, R. J., Rossi, G., Choi, T. K. and Nisonoff, A., X-ray Crystallographic Study of the Fab Fragments from Two Human Myeloma Proteins . . . 223

Low, B. W. and Chen, C. C. H., Monoclinic Insulin Crystals . . . 227

Poglazov, B. F. and Nikolskaya, T. I., Self-assembly of the Protein of Bacteriophage T2 Tail Cores. 231

Woese, C., Models for the Evolution of Codon Assignments . . . 235

Erratum 241

Number 2, 28 July 1969

Ledoux, L. and Huart, R., Fate of Exogenous Bacterial Deoxyribonucleic Acids in Barley Seedlings 243

Wang, J. C., Degree of Superhelicity of Covalently Closed Cyclic DNA's from *Escherichia coli* 263

Rudland, P. S. and Dube, S. K., Specific Interaction of an Initiator tRNA Fragment with 30 s Ribosomal Subunits 273

Englesberg, E., Sheppard, D., Squires, C. and Meronk, F., Jr., An Analysis of "Revertants" of a Deletion Mutant in the *C* gene of the L-Arabinose Gene Complex in *Escherichia coli* B/r: Isolation of Initiator Constitutive Mutants (I^c) 281

Sugiura, M., Okamoto, T. and Takanami, M., Starting Nucleotide Sequences of RNA synthesized on the Replicative Form DNA of Coliphage fd . . 299

Vasquez, C., Kleinschmidt, A. K. and Basilico, C., Electron Microscopic Studies of Polyoma DNA released in Protein Monolayers . . . 317

LETTERS TO THE EDITOR:

APIRION, D., SCHLESSINGER, D., PHILLIPS, S. and SYPHERD, P., *Escherichia coli*: Reversion from Streptomycin Dependence, a Mutation in a Specific 30 s Ribosomal Protein 327

WATSON, J. D. and KIDSON, C., Effects of Ionic Conditions on the Structure of Ribosomal RNA Components 331

KEDES, L. H. and STAVY, L., Structural and Functional Identity of Ribosomes from Eggs and Embryos of Sea Urchins 337

AVEY, H. P. and SHALL, S., Crystallographic Analysis of *N*-Acetyl-DL-Homocysteinyl Ribonuclease Derivatives 341

ROWE, A. J. and WEITZMAN, P. D. J., Allosteric Changes in Citrate Synthase observed by Electron Microscopy 345

WEIL, J., Reciprocal and Non-reciprocal Recombination in Bacteriophage λ 351

NASH, H. A. and MERRIL, C. R., DNA of Biotin-transducing λ Bacteriophage 357

NUMBER 3, 14 AUGUST 1969

SAKORE, T. D., TAVALE, S. S. and SOBELL, H. M., Base-pairing Configurations between Purines and Pyrimidines in the Solid State. I. Crystal and Molecular Structure of a 1:2 Purine–Pyrimidine Hydrogen-bonded Complex: 9-Ethyladenine–1-Methyl-5-iodouracil 361

TAVALE, S. S., SAKORE, T. D. and SOBELL, H. M., Base-pairing Configurations between Purines and Pyrimidines in the Solid State. II. Crystal and Molecular Structure of 9-Ethyl-8-bromoadenine–1-Methyl-5-bromouracil . 375

SAKORE, T. D., SOBELL, H. M., MAZZA, F. and KARTHA, G., Base-pairing Configurations between Purines and Pyrimidines in the Solid State. III. Crystal and Molecular Structure of two 2:1 Hydrogen-bonded Complexes, 1-Methylthymine:9-Ethyl-2,6-diaminopurine and 1-Methyl-5-iodouracil: 9-Ethyl-2,6-diaminopurine 385

MAZZA, F., SOBELL, H. M. and KARTHA, G., Base-pairing Configurations between Purines and Pyrimidines in the Solid State. IV. Crystal and Molecular Structure of two 1:1 Hydrogen-bonded Complexes, 1-Methyl-5-bromouracil:9-Ethyl-2-aminopurine and 1-Methyl-5-fluorouracil:9-Ethyl-2-aminopurine 407

KEITH, H. D., PADDEN, F. J., JR. and GIANNONI, G., Crystal Structures of β-Poly-L-glutamic Acid and its Alkaline Earth Salts 423

OLINS, D. E., Interaction of Lysine-rich Histones and DNA 439

YONATH, A. and TRAUB, W., Polymers of Tripeptides as Collagen Models. IV. Structure Analysis of Poly(L-prolyl-glycyl-L-proline). 461

Appendix—Derivation of Integrated Intensities for Streaks and Spots . 476

Traub, W., Polymers of Tripeptides as Collagen Models. V. An X-ray Study of Poly(L-prolyl-glycyl-glycine) 479

Segal, D. M. and Traub, W., Polymers of Tripeptides as Collagen Models. VI. Synthesis and Structural Investigation of Poly(L-alanyl-L-prolyl-glycine) 487

Segal, D. M., Polymers of Tripeptides as Collagen Models. VII. Synthesis and Solution Properties of Four Collagen-like Polyhexapeptides . . . 497

Segal, D. M., Traub, W. and Yonath, A., Polymers of Tripeptides as Collagen Models. VIII. X-ray Studies of Four Polyhexapeptides 519

Williams, L. S. and Neidhardt, F. C., Synthesis and Inactivation of Aminoacyl-transfer RNA Synthetases during Growth of *Escherichia coli* . 529

Szentkiralyi, E. M. and Oplatka, A., On the Formation and Stability of the Enzymically Active Complexes of Heavy Meromyosin with Actin . . 551

Kiger, J. A., Jr. and Sinsheimer, R. L., Vegetative Lambda DNA. V. Evidence Concerning Single-strand Elongation 567

Salomon, R., Kaye, A. M. and Herzberg, M., Mouse Nuclear Satellite DNA: 5-Methylcytosine Content, Pyrimidine Isoplith Distribution and Electron Microscopic Appearance 581

Lindahl, L. and Forchhammer, J., Evidence for Reduced Breakdown of Messenger RNA during Blocked Transcription or Translation in *Escherichia coli* 593

Székely, M. and Sanger, F., Use of Polynucleotide Kinase in Fingerprinting Non-radioactive Nucleic Acids 607

Nagley, P. and Wake, R. G., Effect of 5-Bromouracil on the Pattern of Deoxyribonucleic Acid Replication in Germinating *Bacillus subtilis* Spores 619

Ray, D. S., Replication of Bacteriophage M13. II. The Role of Replicative Forms in Single-strand Synthesis. 631

Letter to the Editor

Ray, D. S. and Schekman, R. W., Replication of Bacteriophage M13. III. Identification of the Intracellular Single-stranded DNA . . 645

Announcement 649

Index to Volume 43

N-Acetyl-DL-homocysteinyl ribonuclease derivatives, crystallographic analysis, 341
Actin, formation and stability of the enzymically active complexes with heavy meromyosin, 551
Allosteric changes in citrate synthase observed by electron microscopy, 345
Antibodies specific for the purine and pyrimidine nucleosides, effect on *in vitro* priming ability of DNA, 41
APIRION, D., SCHLESSINGER, D., PHILLIPS, S. & SYPHERD, P., *Escherichia coli*: reversion from streptomycin dependence, a mutation in a specific 30 s ribosomal protein, 327
L-Arabinose gene complex in *Escherichia coli* B/r, "revertants" of a deletion mutant in the *C* gene to give initiator constitutive mutants (I^c), 281
Arginine-transfer reaction characteristics in rabbit liver cytoplasm, 163
AVEY, H. P. & SHALL, S., Crystallographic analysis of *N*-acetyl-DL-homocysteinyl ribonuclease derivatives, 341
— (see also HUMPHREY, R. L.), 223

Bacillus subtilis, effect of 5-bromouracil on the pattern of DNA replication in germinating spores, 619
Bacteriophage, DNA of biotin-transducing λ bacteriophage, 357
— fr, role of RNA in the assembly process, 191
— λ, superhelical turns of covalently closed DNA, variation of the average rotation angle, 25
— λ, vegetative DNA, evidence concerning single-strand elongation, 567
— λ, reciprocal and non-reciprocal recombination, 351
— M13, identification of the intracellular single-stranded DNA during replication, 645
— —, the role or replicative forms in single-stranded DNA synthesis, 631
— T2, self-assembly of the tail core protein, 231
Barley seedlings, fate of exogenous bacterial DNA's, 243
BASILICO, C. (see also VASQUEZ, C.), 317
BECKA, L. N. (see also HUMPHREY, R. L.), 223
BECKWITH, J. R. (see also REZNIKOFF, W. S.), 201, 215
BEISER, S. M. (see also WALLACE, S. S.), 41
BHORJEE, J. S., FOWLER, A. V. & ZABIN, I., Biochemical evidence that the operator locus is distinct from the *Z* gene in the *lac* operon of *Escherichia coli*, 219
Biotin-transducing λ bacteriophage DNA, 357
BREMER, H. & MUELLER, K., Substrate requirements for initiation and chain growth of enzymically synthesized RNA, 109
— (see also MUELLER, K.), 89, 105
5-Bromouracil, effect on the pattern of DNA replication in germinating spores of *Bacillus subtilis*, 617

Cell division and DNA replication following a shift to a richer medium, 1
ČERNÁ, J. (see also RYCHLÍK, I.), 13
Chain termination and suppression *in vitro* by the UGA codon, 177
CHEN, C. C. H. (see also LOW, B. W.), 227
CHLÁDEK, S. (see also RYCHLÍK, I.), 13
CHOI, T. K. (see also HUMPHREY, R. L.), 223
Codon assignment evolution, models, 235
—, UGA, *in vitro* chain termination and suppression, 177
Codons, nonsense, identification in yeast, 71
Coliphage fd DNA replicative form, the starting nucleotide sequences of RNA synthesized, 299
Collagen-like polyhexapeptides, X-ray studies, 519
— — —, synthesis and solution properties, 497
— — polymers, an X-ray study of poly(L-prolyl-glycyl-glycine), 479
— — —, structure analysis of poly(L-prolyl-glycyl-L-proline), 461
— — —, synthesis and structural investigation of poly(L-alanyl-L-prolyl-glycine), 487
COOPER, S., Cell division and DNA replication following a shift to a richer medium, 1
Crystallographic analysis of *N*-acetyl-DL-homocysteinyl ribonuclease derivatives, 341

Crystal and molecular structure of a 1 : 2 hydrogen-bonded complex, 9-ethyladenine : 1-methyl-5-iodouracil, 361
— — — — — — two 1 : 1 hydrogen-bonded complexes, 1-methyl-5-bromouracil : 9-ethyl-2-aminopurine and 1-methyl-5-flourouracil : 9-ethyl-2-aminopurine, 407
— — — — — — — 2 : 1 hydrogen-bonded complexes, 1-methylthymine : 9-ethyl-2,6-diaminopurine and 1-methyl-5-iodouracil : 9-ethyl-2,6-diaminopurine, 385
— — — — — 9-ethyl-8-bromoadenine : 9-ethyl-8-bromohypoxanthine, 77
— — — — — 9-ethyl-8-bromoadenine : 1-methyl-5-bromouracil, 375
— structures of β-poly-L-glutamic acid and its alkaline earth salts, 423
Crystals of insulin, monoclinic, 227

Deoxyribonucleic acid, degree of superhelicity of covalently closed cyclic DNA's from *Escherichia coli*, 263
— —, electron microscopic studies of polyoma DNA release in protein monolayers, 317
— —, evidence concerning single-strand elongation in vegetative λ DNA, 567
— —, identification of single-stranded intracellular DNA during replication of bacteriophage M13, 645
— — interaction with lysine-rich histones, 439
— —, *in vitro* effect of antibodies specific for purine and pyrimidine nucleosides on priming ability, 41
— — of biotin-transducing λ bacteriophage, 357
— — — mouse nuclear satellite, 5-methylcytosine content, pyrimidine isoplith distribution and electron microscopic appearance, 581
— — replication following a shift to a richer medium, 1
— — replication pattern in germinating *Bacillus subtilis* spores, effect of 5-bromouracil, 619
— — replicative form, the starting nucleotide sequences of RNA synthesized in coliphage fd, 299
— —, the role of replicative forms in single-strand synthesis in bacteriophage M13, 631
— —, variation of the average rotation angle of the helix and superhelical turns of covalently closed cyclic λ DNA, 25
— acids, fate of exogenous bacterial DNA's in barley seedling, 243
Dinitrophenol treatment without tryptophan, intragenic initiations of transcription of the tryptophan operon in *Escherichia coli*, 51
DUBE, S. K. (see also RUDLAND, P.S.), 273

Electron microscopic appearance of mouse nuclear satellite DNA, 581
— — studies of polyoma DNA released in protein monolayers, 317
— microscopy used to observe allosteric changes in citrate synthetase, 345
ENGLESBERG, E., SHEPPARD, D., SQUIRES, C. & MERONK, F., JR., An analysis of "revertants" of a deletion mutant in the *C* gene of the L-arabinose gene complex in *Escherichia coli* B/r: isolation of initiator constitutive mutants (I^c), 281
Enzymically synthesized RNA, heterogeneous initiation and termination, 89, 105
— — —, substrate requirements for initiation and chain growth, 109
ERLANGER, B. F. (see also WALLACE, S. S.), 41
Escherichia coli B/r, an analysis of "revertants" of a deletion mutant in the *C* gene of the L-arabinose gene complex, isolation of initiator constitutive mutants (I^c), 281
— —, biochemical evidence that the operator locus is distinct from the *z* gene in the *lac operon*, 219
— —, degree of superhelicity of covalently closed cyclic DNA's, 263
— —, evidence for reduced breakdown of mRNA during blocked transcription or translation, 593
— —, genetic evidence that the operator locus is distinct from the *z* gene in the *lac* operon, 215
— —, intragenic initiations of transcription of the tryptophan operon following dinitrophenol treatment without tryptophan, 51
— —, reversion from streptomycin dependence, a mutation in a specific 30 s ribosomal protein, 327
— —, synthesis and inactivation of aminoacyl-tRNA synthetases during growth, 529
9-Ethyl-8-bromoadenine : 1-methyl-5-bromouracil, crystal and molecular structure, 375
9-Ethyl-8-bromoadenine : 9-ethyl-8-bromohypoxanthine, a hydrogen-bonded complex, crystal and molecular structure, 77
9-Ethyladenine : 1-methyl-5-iodouracil, a 1 : 2 hydrogen-bonded complex, crystal and molecular structure, 361
Evolution of codon assignments, models, 235

fd (see Coliphage)
fr (see Bacteriophage)
Fab fragments from two human myeloma proteins, X-ray crystallographic study, 223

FORCHHAMMER, J. (see also LINDAHL, L.), 593
FOWLER, A. V. (see also BHORJEE, J. S.), 219

GIANNONI, G. (see also KEITH, H. D.), 423

Haemoglobin synthesis control, rate of translation of the mRNA for the α and β chains, 123
HALADOVÁ, Z. (see also RYCHLÍK, I.), 13
HAWTHORNE, D.C., Identification of nonsense codons in yeast, 71
HERZBERG, M. (see also SALOMON, R.), 581
Histones, lysine-rich, interaction with DNA, 439
HOHN, T., Role of RNA in the assembly process of bacteriophage fr, 191
HORINISHI, H. (see also SOFFER, R. L.), 163
HUART, R. (see also LEDOUX, L.), 243
HUMPHREY, R. L., AVEY, H. P., BECKA, L. N., POLJAK, R. J., ROSSI, G., CHOI, T. K. & NISONOFF A., X-ray crystallographic study of the Fab fragments from two human myeloma proteins, 223
HUNT, T., HUNTER, T. & MUNRO, A., Control of haemoglobin synthesis: rate of translation of the mRNA for the α- and β-chains, 123
HUNTER, T. (see also HUNT, T.), 123
Hybrid 80 s particles, active, formed from subunits of rat, rabbit and protozoan (*Tetrahymena pyriformis*) ribosomes, 151

IMAMOTO, F., Intragenic initiations of transcription of the tryptophan operon in *Escherichia coli* following dinitrophenol treatment without tryptophan, 51
Immunoglobulin, X-ray crystallographic study of the Fab fragments from two human myeloma proteins, 223
Initiation and chain growth of enzymically synthesized RNA, substrate requirements, 109
Initiator constitutive mutants (I^c) in *Escherichia coli*: from a study of the L-arabinose gene complex, 281
– tRNA fragment, specific interaction with 30 s ribosomal subunits, 273
Insulin, monoclinic crystals, 227

KARTHA G. (see also MAZZA, F.), 407
– (see also SAKORE, T. D.) 385
KAYE, A. M. (see also SALOMON, R.), 581
KEDES, L. H. & STAVY, L., Structural and functional identity of ribosomes from eggs and embryos of sea urchins, 337
KEITH, H. D., PADDEN, F. J. & GIANNONI, G., Crystal structures of β-poly-L-glutamic acid and its alkaline earth salts, 423
KIDSON, C. (see also WATSON, J. D.), 331
KIGER, J. A. & SINSHEIMER, R. L., Vegetative lambda DNA. V. Evidence concerning single-strand elongation, 567
KLEINSCHMIDT, A. K. (see also VASQUEZ, C.), 317

λ (see Bacteriophage)
lac operon of *Escherichia coli*, biochemical evidence that the operator locus is distinct from the *z* gene, 219
– — — — —, genetic evidence that the operator locus is distinct from the *z* gene, 215
LEDOUX, L. & HUART, R., Fate of exogenous bacterial deoxyribonucleic acids in barley seedlings, 243
LINDAHL, L. & FORCHHAMMER, J., Evidence for reduced breakdown of messenger RNA during blocked transcription or translation in *Escherichia coli*, 593
LOW, B. W. & CHEN, C. C. H., Monoclinic insulin crystals, 227
LOW, R. B. (see also MARTIN, T. E.), 135
Lysine-rich histones, interaction with DNA, 439

M13 (see Bacteriophage)
MARTIN, T. E., ROLLESTON, F. S., LOW, R. B. & WOOL, I. G., Dissociation and reassociation of skeletal muscle ribosomes, 135
– & WOOL, I. G., Active hybrid 80 s particles formed from subunits of rat, rabbit and protozoan (*Tetrahymena pyriformis*) ribosomes, 151

Mazza, F., Sobell, H. M. & Kartha, G., Base-pairing configurations between purines and pyrimidines in the solid state. IV. Crystal and molecular structure of two 1:1 hydrogen-bonded complexes, 1-methyl-5-bromouracil:9-ethyl-2-aminopurine, and 1-methyl-5-fluorouracil:9-ethyl-2-aminopurine, 407
— (see also Sakore, T. D.), 385
Meromyosin, heavy, formation and stability of the enzymically active complexes with actin, 557
Meronk, F., Jr., (see also Englesberg, E.), 281
Merril, C. R. (see also Nash, H. A.), 357
Messenger RNA evidence for reduced breakdown during blocked transcription or translation in *Escherichia coli*, 593
— —, rate of translation for the α- and β-chains during haemoglobin synthesis, 123
1-Methyl-5-bromouracil:9-ethyl-2-aminopurine, a 1:1 hydrogen-bonded complex, crystal and molecular structure, 407
1-Methyl-5-fluorouracil:9-ethyl-2-aminopurine, a 1:1 hydrogen-bonded complex, crystal and molecular structure, 407
1-Methyl-5-iodouracil:9-ethyl-2,6-diaminopurine, a 2:1 hydrogen-bonded complex, crystal and molecular structure, 385
5-Methylcytosine content of mouse nuclear satellite DNA, 581
1-Methylthymine:9-ethyl-2,6-diaminopurine, a 2:1 hydrogen-bonded complex, crystal and molecular structure, 385
Miller, J. H. (see also Reznikoff, W. S.), 201
Model, P., Webster, R. E. & Zinder, N. D., The UGA codon *in vitro* chain termination and suppression, 177
Mueller, K. & Bremer, H., *Appendix*: Termination and length of RNA molecules at different reaction times, 105
— & —, Heterogeneous initiation and termination of enzymically synthesized ribonucleic acid, 89
— (see also Bremer, H.), 109
Munro, A. (see also Hunt, T.), 123
Muscle ribosomes, skeletal, dissociation and reassociation, 135
Myeloma protein, human, X-ray crystallographic study of the Fab fragment, 223

Nagley, P. & Wake, R. G., Effect of 5-bromouracil on the pattern of DNA replication in germinating *Bacillus subtilis* spores, 619
Nash, H. A. & Merril, C. R., DNA of biotin-transducing λ bacteriophage, 357
Neidhardt, F. C. (see also Williams, L. S.), 529
Nikolskaya, T. I. (see also Poglazov, B. F.), 231
Nisonoff, A. (see also Humphrey, R. L.), 223
Nonsense codons, indentification in yeast, 71
Nucleic acids, use of polynucleotide kinase in fingerprinting non-radioactive nucleic acids, 605

Okamoto, T. (see also Sugiura, M.), 299
Olins, D. E., Interaction of lysine-rich histones and DNA, 439
Operator locus, biochemical evidence that it is distinct from the *z* gene in the *lac* operon of *Escherichia coli*, 219
— —, genetic evidence that it is distinct from the *z* gene in the *lac* operon of *Escherichia coli*, 215
Oplatka, A. (see also Szentkiralyi, E. M.), 551

Padden, F. J. (see also Keith, H. D.), 423
Peptidyl transferase, ribosomal, 2′ (3′)-*O*-aminoacyl nucleosides as acceptors of the peptide chain on the amino acid site, 13
Phillips, S. (see also Apirion, D.), 327
Poglazov, B. F. & Nikolskaya, T. I., Self-assembly of the protein of bacteriophage T2 tail cores, 231
Poljak, R. J. (see also Humphrey, R. L.), 223
Poly (L-alanyl-L-prolyl-glycine) as a collagen model, synthesis and structural studies, 487
— (L-prolyl-glycyl-glycine) as a collagen model, an X-ray study, 479
— (L-prolyl-glycyl-L-proline) as a collagen model, structure analysis, 461
β-Poly-L-glutamic acid and its alkaline earth salts, crystal structures, 423
Polyhexapeptides, collagen-like, synthesis and solution properties, 497
—, — —, X-ray studies, 519
Polynucleotide kinase, use in fingerprinting non-radioactive nucleic acids, 607

Polyoma DNA released in protein monolayers studied by electron microscopy, 317
Protozoan, *Tetrahymena pyriformis*, ribosomes, active hybrid 80 s particles formed with subunits from rat and rabbit ribosomes, 151
Purines and pyrimidines, crystal and molecular structure of a 1:2 hydrogen-bonded complex, 9-ethyladenine:1-methyl-5-iodouracil, 361
— — —, — — — — 9-ethyl-8-bromoadenine:1-methyl-5-bromouracil, 375
— — —, — — — — two 1:1 hydrogen-bonded complexes, 1-methyl-5-bromouracil:9-ethyl-2-aminopurine and 1-methyl-5-fluorouracil:9-ethyl-2-aminopurine, 407
— — —, — — — — — two 2:1 hydrogen-bonded complexes, 1-methylthymine:9-ethyl-2, 6-diaminopurine and 1-methyl-5-iodouracil:9-ethyl-2, 6-diaminopurine, 385
— — — nucleoside-specific antibodies, effect on *in vitro* priming ability of DNA, 41
Pyrimidine isoplith distribution of mouse nuclear satellite DNA, 581

Rabbit liver cytoplasm, characteristics of the arginine-transfer reaction, 163
Ray, D. S., Replication of bacteriophage M13. II. The role of replicative forms in single-strand synthesis, 631
— & Schekman, R. W., Replication of bacteriophage M13. III. Identification of the intracellular single-stranded DNA, 645
Replication of DNA following a shift to a richer medium, 1
Repressor action, mechanism, 201
Reznikoff, W. S. & Beckwith, J. R., Genetic evidence that the operator locus is distinct from the *z* gene in the *lac* operon of *Escherichia coli*, 215
—, Miller, J. H., Scaife, J. G. & Beckwith, J. R., A mechanism for repressor action, 201
Ribonuclease, *N*-acetyl-DL-homocysteinyl derivatives, crystallographic analysis, 341
Ribonucleic acid, aminoacyl-transfer synthetases, synthesis and inactivation during growth of *Escherichia coli*, 529
— —, enzymically synthesized, heterogeneous initiation and termination, 89, 105
— —, — —, substrate requirements for initiation and chain growth, 109
— —, messenger, evidence for reduced breakdown during blocked transcription or translation in *Escherichia coli*, 593
— —, —, rate of translation for the α- and β-chains during haemoglobin synthesis, 123
— —, ribosomal components, effects of ionic conditions on structure, 331
— —, role in the assembly process of bacteriophage fr, 191
— —, specific interaction of an initiator tRNA fragment with 30 s ribosomal subunits, 273
— — synthesized on the replicative form DNA of coliphage fd, the starting nucleotide sequences, 299
Ribosomal 30 s subunits, specific interaction with an initiator tRNA fragment, 273
— peptidyl transferase substrate specificity, 2′ (3′)-*O*-aminoacyl nucleosides as acceptors of the peptide chain on the amino acid site, 13
— protein, 30 s, a specific mutation reverses streptomycin dependence in *Escherichia coli*, 327
— RNA components, effects of ionic conditions on structure, 331
Ribosome hybrid 80 s particles, from subunits of rat, rabbit and protozoa (*Tetrahymena pyriformis*) ribosomes, 151
Ribosomes from eggs and embryos of sea urchins, structural and functional identity of ribosomes, 337
— of skeletal muscle, dissociation and reassociation, 135
Rolleston, F. S. (see also Martin, T. E.), 135
Rossi, G. (see also Humphrey, R. L.), 223
Rowe, A. J. & Weitzman, P. D. J., Allosteric changes in citrate synthase observed by electron microscopy, 345
Rudland, P. S. & Dube, S. K., Specific interaction of an initiator tRNA fragment with 30 s ribosomal subunits, 273
Rychlík, I., Černá, J., Chládek, S., Zemlička, J. & Haladová, Z., Substrate specificity of ribosomal peptidyl transferase: 2′ (3′)-*O*-aminoacyl nucleosides as acceptors of the peptide chain on the amino acid site, 13

Sakore, T. D. & Sobell, H. M., The crystal and molecular structure of a hydrogen-bonded complex containing adenine and hypoxanthine derivatives: 9-ethyl-8-bromoadenine: 9-ethyl-8-bromohypoxanthine, 77
—, —, Mazza, F. & Kartha, G., Base-pairing configurations between purines and pyrimidines in the solid state. III. Crystal and molecular structure of two 2:1 hydrogen-bonded complexes, 1-methylthymine:9-ethyl-2,6-diaminopurine and 1-methyl-5-iodouracil:9-ethyl-2,6-diaminopurine, 385

Sakore, T. D., Tavale, S. S. & Sobell, H. M., Base-pairing configurations between purines and pyrimidines in the solid state. I. The crystal and molecular structure of a 1:2 purine–pyrimidine hydrogen-bonded complex; 9-ethyladenine : 1-methyl-5-iodouracil, 361
— (see also Tavale, S. S.), 375
Salomon, R., Kaye, A. M. & Herzberg, M., Mouse nuclear satellite DNA : 5-methylcytosine content, pyrimidine isoplith distribution and electron microscopic appearance, 581
Sanger, F. (see also Szekely, M.), 607
Scaife, J. G. (see also Reznikoff, W. S.), 201
Schekman, R. W. (see also Ray, D. S.), 645
Schlessinger, D. (see also Apirion, D.), 327
Sea urchin eggs and embryos, structural and functional identity of ribosomes, 337
Segal, D. M., Polymers of tripeptides as collagen models. VII. Synthesis and solution properties of four collagen-like polyhexapeptides, 497
— & Traub, W., Polymers of tripeptides as collagen models. VI. Synthesis and structural investigation of poly(L-alanyl-L-prolyl-glycine), 487
—, — & Yonath, A., Polymers of tripeptides as collagen models. VIII. X-ray studies of four polyhexapeptides, 519
Shall, S. (see also Avey, H. P.), 341
Shepherd, D. (see also Englesberg, E.), 281
Sinsheimer, R. L. (see also Kiger, J. A.), 567
Sobell, H. M. (see also Mazza, F.), 407
— (see also Sakore, T. D.), 77, 361, 385
— (see also Tavale, S. S.), 375
Soffer, R. L. & Horinishi, H., Enzymic modification of proteins. General characteristics of the arginine-transfer reaction in rabbit liver cytoplasm, 163
Squires, C. (see also Englesberg, E.), 281
Stavy, L. (see also Kedes, L. H.), 337
Streptomycin dependence, reversion in *Escherichia coli*, a mutation in a specific 30 s ribosomal protein, 327
Structural and functional identity of ribosomes from eggs and embryos of sea urchins, 337
Sugiura, M., Okamoto, T. & Takanami, M., Starting nucleotide sequences of RNA synthesized on the replicative form DNA of coliphage fd, 299
Suppression and chain termination *in vitro* by the UGA codon, 177
Synthase citrate, allosteric changes observed by electron microscopy, 345
Sypherd, P. (see also Apirion, D.), 327
Szekely, M. & Sanger, F., Use of polynucleotide kinase in fingerprinting non-radioactive nucleic acids, 607
Szentkiralyi, E. M. & Oplatka, A., On the formation and stability of the enzymically active complexes of heavy meromyosin with actin, 551

T2 (see Bacteriophage)
Takanami, M. (see also Sugiura, M.), 299
Tavale, S. S., Sakore, T. D. & Sobell, H. M., Base-pairing configurations between purines and pyrimidines in the solid state. II. Crystal and molecular structure of 9-ethyl-8-bromoadenine: 1-methyl-5-bromouracil, 375
— (see also Sakore, T. D.), 361
Tetrahymena pyriformis (see Protozoan)
Transfer RNA, specific interaction of an initiator tRNA fragment with 30 s ribosomal subunits, 273
— —, synthesis and inactivation of aminoacyl-tRNA synthetases during growth of *Escherichia coli*, 529
Traub, W., Polymers of tripeptides as collagen models. V. An X-ray study of poly(L-prolyl-glycyl-glycine), 479
— (see also Segal, D. M.), 487, 519
— (see also Yonath, A.), 461, 477
Tryptophan operon transcription in *Escherichia coli*, intragenic initiations following dinitrophenol treatment without tryptophan, 51

Vasquez, C., Kleinschmidt, A. K. & Basilico, C., Electron microscopic studies of polyoma DNA released in protein monolayers, 317

Wake, R. G. (see also Nagley, P.), 619
Wallace, S. S., Erlanger, B. F. & Beiser, S. M., Antibodies specific for purine and pyrimidine nucleosides: effect on *in vitro* priming ability of DNA, 41

Wang, J. C., Degree of superhelicity of covalently closed cyclic DNA's from *Escherichia coli*, 263
—, Variation of the average rotation angle of the DNA helix and the superhelical turns of covalently closed cyclic λ DNA, 25
Watson, J. D. & Kidson, C., Effects of ionic conditions on the structure of ribosomal RNA components, 331
Webster, R. E. (see also Model, P.), 177
Weil, J., Reciprocal and non-reciprocal recombination in phage λ, 351
Weitzman, P. D. J. (see also Rowe, A. J.), 345
Williams, L. S. & Neidhardt, F. C., Synthesis and inactivation of aminoacyl-tRNA synthetases during growth of *Escherichia coli*, 529
Woese, C. R., Models for the evolution of codon assignments, 235
Wool, I. G. (see also Martin, T. E.), 135, 151

X-ray crystallographic study of Fab fragments from two human myeloma proteins, 223
— study of four polyhexapeptides as a collagen model, 519
— — — poly(L-prolyl-glycyl-glycine) as a collagen model, 479

Yonath, A. & Traub, W., *Appendix* : Derivation of integrated intensities for streaks and spots, 475
— & —, Polymers of tripeptides as collagen models. IV. Structure analysis of poly(L-prolyl-glycyl-L-proline), 461
— (see also Segal, D. M.), 519
Yeast, identification of nonsense codons, 71

Zabin, I. (see also Bhorjee, J. S.), 219
Zemlička, J. (see also Rychlík, I.), 13
Zinder, N. D. (see also Model, P.), 177